



LACK OF STABLE INITIATION FACTOR 3 (IF-3) BINDING TO DIMERS OF THE 30 S RIBOSOMAL SUBUNITS*

Claudio GUALERZI[†], Matthias R. WABL and Cynthia L. PON^{**}
*Max-Planck-Institut für Molekulare Genetik, Abteilung Wittmann,
Innestrasse 63–73, 1, Berlin 33 (Dahlem) Germany*

Received 25 June 1973

1. Introduction

It is known that initiation factor 3 (IF-3) is endowed with ribosome dissociation factor (DF) activity [1, 2]. Recent studies have shown that IF-3 acts as anti-association factor since the equilibrium between 70 S ribosomes and 50 S and 30 S ribosomal subunits is shifted in favour of the subunits by the binding of IF-3 to free 30 S particles [3–5]. The finding that IF-3 does not *stably* bind to the 70 S monomers is consistent with this scheme [6–8]. The 30 S–IF-3 complex is dissociated during the physiological formation of the 70 S initiation complex upon addition of 50 S ribosomal subunits in the presence of fMet-tRNA, or during the formation of 70 S couples when 30 S and 50 S particles are 'forced together' at high (15 mM) Mg^{2+} concentration [6]. Therefore, an antagonism between IF-3 and 50 S particles for the binding to the 30 S ribosomal subunits seems to exist.

In the present paper we present evidence that the IF-3 binding capacity of the 30 S ribosomal particles is weakened or lost when these particles are engaged in the formation of 30 S–30 S dimers and suggest that an analogy may exist between 30 S–30 S and 30 S–50 S interactions.

2. Materials and methods

The preparation of *E. coli* (MRE 600) 30 S ribosomal subunits will be described in detail in a forthcoming paper [9]. The purification of IF-3 was carried out as previously reported [10]. Labelling *in vitro* of IF-3 was carried out by reductive alkylation [11] as previously described [7]. In a typical preparation, purified (electrophoretically homogeneous) IF-3 was concentrated to approximately 3.0 mg/ml and exhaustively dialyzed versus 0.1 M sodium borate buffer (pH 9.0) containing 0.2 M KCl and 5 mM 2-mercaptoethanol. ^{14}C -formaldehyde (57.6 mCi/mmol) was then added (0.25 mCi/1.65 mg of protein) and the mixture incubated for 30 sec in an ice-bath. After incubation sodium borohydride was added until a 2-fold molar excess of sodium borohydride over formaldehyde was reached. The solution was then dialyzed exhaustively against several changes of a buffer containing 20 mM Tris–HCl (pH 7.4); 0.5 mM dithiothreitol (DTT); 1 mM EDTA; 0.2 M NH_4Cl and 5% glycerol.

For the binding of IF-3 to ribosomes, 4 μg of [^{14}C]H₃–IF-3 (21,000 cpm/ μg) were incubated with 0.30 A_{260} units of 30 S ribosomal subunits in 0.4 ml of buffer containing 10 mM Tris–HCl (pH 7.5); 100 mM NH_4Cl ; 5 mM 2-mercaptoethanol, either 1 or 10 mM (cf. fig. 1) $Mg (Ac)_2$ and 0.2 mM GTP. After 10 min at 37°C, each incubation mixture was transferred onto the top of a 10–30% (w/v) sucrose density gradient made in the above buffer (either 1 or 10 mM $Mg (Ac)_2$) and centrifuged for 3 hr at 48 000 rpm at 2°C in a SW 50.1 rotor. Each gradient was pumped through the flow-cell of a Zeiss spectrophotometer and fractionated directly into counting vials. The radioactivity of each fraction

[†] Address correspondence to: Dr. C. Gualerzi, Institut de Biologie Moléculaire, Faculté des Sciences de Paris, 2, Place Jussieu, Paris V.

* Paper no. 62 on 'Ribosomal Proteins'. Preceding paper is ref. 19.

** On leave from the Dept. of Biological Sciences, Hunter College of the City University of New York, N.Y. 10021, USA.




Fig. 1. Binding of [^{14}C]H $_3$ -IF-3 to 30 S ribosomal subunits. The binding reaction and the subsequent centrifugation were carried out as described in Materials and methods (●····●····●) [^{14}C] radioactivity; (—) A_{260} . Sedimentation was from right to left.

some preparation (age, number of cycles of freezing and thawing) and on the Mg^{2+} concentration of the resuspending medium.

The binding of IF-3 to a preparation of 30 S particles which displayed a marked tendency to dimerize at high (10–20 mM) Mg^{2+} is shown in fig. 1. As seen in the figure, when the binding of radioactive IF-3 to ribosomes and the subsequent centrifugation through sucrose density gradient were carried out in a buffer containing 10 mM Mg^{2+} , a discrete A_{260} peak of 30 S-dimers sedimented faster than the 30 S peak. All the IF-3 radioactivity was associated with the 30 S peak and no IF-3 sedimented with the 30 S-dimers under these conditions. If the binding of IF-3 and the subsequent centrifugation were carried out at 1 mM Mg^{2+} , however, the A_{260} dimer peak was reduced to a shoulder, while the 30 S peak showed a corresponding increase. In addition, the increase in the A_{260} of the 30 S particles resulted in a concomitant increase in the amount of radioactive IF-3 sedimenting in the 30 S region, while the remaining fast-moving shoulder was still free of radioactivity. The possibility that the increase in radioactive IF-3 sedimenting in the 30 S region at 1 mM Mg^{2+} could merely reflect an increased binding affinity of the non-dimerizing 30 S subunits for IF-3, can be ruled out for the following reasons: a) in the experiment shown in fig. 1 a large molar excess of IF-3 over 30 S ribosomal subunits has been used so that each ribosomal subunit is 'saturated' with one molecule of IF-3; b) changes in the Mg^{2+} concentration between 1 and 15 mM do not influence the affinity of the 30 S

was determined after addition of 10 ml of Bray's solution.

3. Results

The formation of dimers of the 30 S ribosomal subunits of prokaryotic ribosomes is a known, although poorly understood, phenomenon [12, 13]. In our experience, the dimerization of the 30 S ribosomal subunits of *E. coli* is partially dependent on the ribo-




Fig. 2. Electron micrographs of negatively stained 30 S-30 S-dimers. The 30 S-dimers were dialyzed against 10 mM Tris-HCl (pH 7.6); 20 mM $\text{Mg}(\text{Ac})_2$; 60 mM NH_4Cl and 6 mM 2-mercaptoethanol and then incubated 10 min at 37°C with 1% uranyl oxalate, pH 6.8. Magnification 400 000X. a, 1–4, b, 1–2 typical dimer formation; b, 3–4 atypical dimer formation.




Fig. 3. Electron micrographs of negatively stained 70 S ribosomes. The 70 S monomers, in 10 mM Tris-HCl (pH 7.6); 10 mM Mg (Ac)₂; 60 mM NH₄Cl and 6 mM 2-mercaptoethanol, were stained with 0.5% uranyl acetate. Magnification 325 000X. The arrows indicate the 30 S subunits.

ribosomal subunits for IF-3 [6–8, 14]; c) when the particles sedimenting in the 30 S-dimer peak are isolated and dissociated at low Mg²⁺, the 30 S ribosomal subunits so obtained are capable of binding radioactive IF-3 (not shown).

Recent studies have shown that the 30 S ribosomal subunits are ellipsoidal and sometimes curved particles of uniform size bearing a cleft in an asymmetrical position perpendicular to the long axis of the particle. This cleft, which divides the 30 S particle into two unequal lobes, is more pronounced on one side of the particle [15, 16].

Eight electron micrographs of 30 S-dimers are presented in fig. 2. Typical 30 S-dimers (over 90% of the dimers examined) appear to be particles of relatively homogeneous size and shape with the two 30 S subunits facing each other on the side where the cleft is more pronounced (fig. 2a 1–4, b 1–2). Fig. 2,b3 shows an atypical dimer in which the 30 S subunits interact 'back to back'. Furthermore, in the majority of the dimers the 30 S subunits seem to face each other in an 'antiparallel' fashion so that the large lobe of one subunit always touches the small lobe of the other (fig. 2 a 1–4, b 1–2). Fig. 2 b4 shows an exceptional case of two subunits aligned parallel to each other. In the majority of the cases, therefore, the formation of 30 S-dimers occurs through the interaction of specific regions of the 30 S particles and not through random contact between the two particles.

Examination of the electron micrographs of 70 S couples (fig. 3) seems to indicate that the 30 S subunits are in contact with the 50 S subunits through the same

'concave' surface which is responsible for the 30 S–30 S interaction (note the cleft dividing the 30 S subunits which now appears as a slightly eccentric 'hole' at the interface between the 30 S and 50 S subunit) suggesting an analogy between 30 S–30 S and 30 S–50 S interactions.

4. Discussion

The above results do not rule out the possibility of a weak interaction between IF-3 and 30 S-dimers but show that the 30 S ribosomal subunit cannot *stably* bind IF-3 if another ribosomal particle (either 30 S or 50 S) is associated with it. In almost all the 30 S-dimers observed, the 30 S particles are in contact through a specific region of their surface (most likely the same region that interacts with the 50 S subunits in the 70 S couples). It would be tempting, therefore, to speculate that also the ribosomal binding site for IF-3 [17] is localized on this part of the 30 S particles. Another explanation (which does not necessarily exclude the first one) for the lack of *stable* IF-3 binding to 30 S-dimers could be that when the 30 S subunits are in association with another ribosomal particle (30 S or 50 S), they assume a configuration different from that of the free 30 S and not compatible with a *stable* interaction with IF-3. Evidence has been presented that the 30 S particle undergoes a configurational change during association with 50 S subunits [18] and the importance of the configuration of the 30 S particles in the binding of IF-3 has already been pointed out [17]. The ability of IF-3 to induce a

configurational transition of the 30 S subunits will be described in a forthcoming paper [9].

Acknowledgements

We wish to express our gratitude to Professor H.G. Wittmann for his encouragement and constructive discussion, Dr. A.L. Haenni for reviewing the manuscript and Miss Heike Jager for the skilful technical assistance. We are also grateful to the A. von Humboldt Stiftung for a Forshungsstipendium given to one of us (C.G.) during the course of this study.

References

- [1] Sabol, S., Sillero, M.A.G., Iwasaki, K. and Ochoa, S. (1970) *Nature* 228, 1269.
- [2] Subramanian, A.R. and Davis, B.D. (1970) *Nature* 228, 1273.
- [3] Kaempfer, R. (1971) *Proc. Natl. Acad. Sci. U.S.* 68, 2458.
- [4] Noll, M. and Noll, H. (1972) *Nature New Biol.* 238, 225.
- [5] Sabol, S., Meier, D. and Ochoa, S. (1973) *European J. Biochem.* 33, 332.
- [6] Sabol, S. and Ochoa, S. (1971) *Nature New Biol.* 234, 233.
- [7] Pon, C.L., Friedman, S.M. and Gualerzi, C. (1972) *Mol. Gen. Genet.* 116, 192.
- [8] Thibault, J., Chestier, A., Vidal, D. and Gros, F. (1972) *Biochimie* 54, 829.
- [9] Paradies, H.H., Franz, A., Pon, C.L. and Gualerzi, C., in preparation.
- [10] Gualerzi, C., Pon, C.L. and Kaji, A. (1971) *Biochem. Biophys. Res. Commun.* 45, 1312.
- [11] Rice, R.H. and Means, G.E. (1971) *J. Biol. Chem.* 246, 831.
- [12] Gualerzi, C. and Cammarano, P. (1969) *Biochim. Biophys. Acta* 190, 551.
- [13] VanDiggelen, O., Ph.D. Thesis (1973) Dept. of Biochemistry, University of Leiden.
- [14] Bosch, L., Veermer, C., Bloemsma-Jonkman, F. and Talens, J. Abstracts of the 8th Fed. European Biochem. Meet., Amsterdam, 517, 1972.
- [15] Wabl, M.R., Deberer, H.G., Höglund, S. and Ljung, L. (1973) *Cytobiologie* 7, 111.
- [16] Wabl, M.R., Barends, P. and Nanninga, N. (1973) *Cytobiologie* 7, 1.
- [17] Gualerzi, C. and Pon, C.L. (1973) *Biochem. Biophys. Res. Commun.*, 52, 792.
- [18] Sherman, M.I. and Simpson, M.V. (1969) *Cold Spring Harbor Symp. Quant. Biol.* 34, 220.
- [19] Wittmann, H.G. et al. (1973) *Mol. Gen. Genet.*, in press.